



ERO

U.S. Immigration and Customs Enforcement Enforcement and Removal Operations

COVID-19 Pandemic Response Requirements



**U.S. Immigration
and Customs
Enforcement**

Table of Contents

SUMMARY OF CHANGES	3
PURPOSE AND SCOPE.....	5
INTRODUCTION	5
OBJECTIVES	6
COMPLIANCE MEASURES.....	7
CONCEPT OF OPERATIONS.....	8
DEDICATED ICE DETENTION FACILITIES	8
NON-DEDICATED ICE DETENTION FACILITIES.....	10
ALL FACILITIES HOUSING ICE DETAINEES.....	12
PREPAREDNESS	12
PREVENTION	17
MANAGEMENT	20
ATTACHMENTS	25

Summary of Changes

Explanation of Change	Section	Page
<ul style="list-style-type: none"> • <u>Addition</u>: Other symptoms of COVID-19 may include chills, muscle pain, sore throat, and new loss of taste or smell. 	Introduction	6
<ul style="list-style-type: none"> • <u>Reiteration</u>: ICE detention standards require facilities to collaborate and comply with federal, state, and/or local authorities addressing public health issues, including communicable diseases. 	Introduction	6
<ul style="list-style-type: none"> • <u>Reiteration</u>: The ERO PRR establishes mandatory requirements, as well as best practices, for all ICE detention facilities in response to COVID-19. 	Objectives	6
<ul style="list-style-type: none"> • <u>Addition</u>: Compliance measures are added to explain how ICE will ensure ICE detention facilities comply with ERO PRR requirements. 	Compliance Measures	7
<ul style="list-style-type: none"> • <u>Addition</u>: New intakes into ICE custody at dedicated ICE detention facilities will be evaluated within five days to determine if the detainee falls within the populations identified by the CDC as potentially being at higher risk for serious illness from COVID-19 and/or the <i>Fraihat v. ICE</i> subclasses. 	Concept of Operations: Dedicated ICE Detention Facilities	8
<ul style="list-style-type: none"> • <u>Addition</u>: Detainees at ICE dedicated facilities who claim to fall within the populations identified by the CDC as potentially being at higher risk for serious illness from COVID-19 should be evaluated within five days, and the ERO Field Officer Director and Field Medical Coordinator notified within 12 hours of determining whether the detainee meets the criteria. 	Concept of Operations: Dedicated ICE Detention Facilities	8
<ul style="list-style-type: none"> • <u>Addition</u>: The ERO PRR identifies additional populations potentially at higher risk for serious illness from COVID-19 (e.g., severe psychiatric illness, pregnancy, autoimmune diseases, and persons of all ages with a physical or mental impairment that substantially limits one or more major life activities). 	Concept of Operations: Dedicated ICE Detention Facilities & Non-Dedicated ICE Detention Facilities	8-10

<ul style="list-style-type: none"> <u>Addition</u>: The ERO PRR previously suspended volunteer work program (VWP) assignments only for detainees assigned to food service; it now also suspends any VWP assignment that requires interaction between individuals at a distance of less than six feet, and requires any detainee participating in VWP to wear job appropriate PPE during work hours. 	Prevention: Visitation	15
<ul style="list-style-type: none"> <u>Clarification</u>: All facility staff (both medical and correctional) are required to wear PPE when encountering or interacting with a detainee at less than six feet. 	Prevention: Visitation	15
<ul style="list-style-type: none"> <u>Addition</u>: PPE is required whenever staff interact with an individual who appears feverish or ill and/or with respiratory symptoms. 	Prevention: Visitation	15-16
<ul style="list-style-type: none"> <u>Addition</u>: If single isolation rooms are unavailable, individuals with confirmed COVID-19 should be isolated together as a cohort separate from other detainees, including those with pending test results. 	Management: For Suspected or Confirmed COVID-19 Cases	17
<ul style="list-style-type: none"> <u>Addition</u>: Housing of suspected or confirmed COVID-19 cases should maintain separation of groups by common criteria. 	Management: For Suspected or Confirmed COVID-19 Cases	17
<ul style="list-style-type: none"> <u>Addition/Clarification</u>: The ERO PRR updates and clarifies the criteria for discontinuation of transmission-based precautions prior to housing detainees in general population. 	Management: For Suspected or Confirmed COVID-19 Cases	18
<ul style="list-style-type: none"> <u>Addition</u>: Links are provided to CDC recommendations for viral testing and specimen collection, as well as for antibody testing. 	Management: For Suspected or Confirmed COVID-19 Cases	19

PURPOSE AND SCOPE

The U.S. Immigration and Customs Enforcement (ICE) Enforcement and Removal Operations (ERO) Coronavirus Disease 2019 (COVID-19) Pandemic Response Requirements (PRR) sets forth expectations and assists ICE detention facility operators to sustain detention operations while mitigating risk to the safety and wellbeing of detainees, staff, contractors, visitors, and stakeholders due to COVID-19. Consistent with ICE's overall adjustments to its immigration enforcement posture,¹ the ERO PRR builds upon previously issued guidance and sets forth specific mandatory requirements to be adopted by all detention facilities, as well as recommended best practices, to ensure that detainees are appropriately housed and that available mitigation measures are implemented during this unprecedented public health crisis.² The ERO PRR has been developed in consultation with the Centers for Disease Control and Prevention (CDC) and is a dynamic document that will be updated as additional/revised information and best practices become available.

INTRODUCTION

As the CDC has explained:

COVID-19 is a communicable disease caused by a novel (new) coronavirus, SARS-CoV-2, and was first identified as the cause of an outbreak of respiratory illness that began in Wuhan Hubei Province, People's Republic of China (China).

COVID-19 appears to spread easily and sustainably within communities. The virus is thought to transfer primarily by person-to-person contact through respiratory droplets produced when an infected person coughs or sneezes; it may transfer through contact with surfaces or objects contaminated with these droplets. There is also evidence of asymptomatic transmission, in which an individual infected with COVID-19 is capable of spreading the virus to others before exhibiting symptoms. The ease of transmission presents a risk of a surge in hospitalizations for COVID-19, which would reduce available hospital capacity. Such a surge has been identified as a likely contributing factor to the high mortality rate for COVID-19 cases in Italy and China.

Symptoms include fever, cough, and shortness of breath, and typically appear two to fourteen days after exposure. Manifestations of severe disease include severe pneumonia, acute respiratory distress syndrome (ARDS), septic shock, and multi-

¹ See, e.g., Attachment A, U.S. Immigration and Customs Enforcement, Updated ICE statement on COVID-19 (Mar. 18, 2020), <https://www.ice.gov/news/releases/updated-ice-statement-covid-19>.

² On April 20, 2020, the U.S. District Court for the Central District of California issued a preliminary injunction requiring that ICE "issue a performance standard or a supplement to their Pandemic Response Requirements ... defining the minimum acceptable detention conditions for detainees with risk factors." *Fraihat v. ICE*, --- F.Supp.3d ---, 2020 WL 1932570, *29 (C.D. Cal. Apr. 20, 2020). The ERO PRR has accordingly been updated to define the "minimum acceptable detention conditions for detainees with risk factors."

organ failure. According to the World Health Organization, approximately 3.4 percent of reported COVID-19 cases have resulted in death globally. This mortality rate is higher among older adults or those with compromised immune systems. Older adults and people who have severe chronic medical conditions like heart, lung, or kidney disease are also at higher risk for more serious COVID-19 illness. Early data suggest older people are twice as likely to have serious COVID-19 illness.

Notice of Order Under Sections 362 and 365 of the Public Health Service Act Suspending Introduction of Certain Persons From Countries Where a Communicable Disease Exists, 85 Fed. Reg. 17060 (Mar. 26, 2020) (internal citations omitted).

Additionally, other symptoms may include chills, muscle pain, sore throat, and new loss of taste or smell.³ Given the seriousness and pervasiveness of COVID-19, ICE is taking necessary and prompt measures. ICE is providing guidance on the minimum measures required for facilities housing ICE detainees to implement to ensure consistent practices throughout its detention operations and the provision of medical care across the full spectrum of detention facilities to mitigate the spread of COVID-19. The ICE detention standards applicable to all facilities used to house ICE detainees have long required that each such facility have written plans that address the management of infectious and communicable diseases, including, but not limited to, testing, isolation, prevention, treatment, and education. Those requirements include reporting and collaboration with local or state health departments in accordance with state and local laws and recommendations.⁴

The Performance-Based National Detention Standards (PBNDS) 2008 and 2011 both require facilities to “comply with current and future plans implemented by federal, state or local authorities addressing specific public health issues including communicable disease reporting requirements.” The 2019 National Detention Standards (NDS) similarly require “collaboration with local or state health departments in accordance with state and local laws and recommendations.” The measures set forth in the ERO PRR allow ICE personnel and detention providers to properly discharge their obligations under those standards in light of the unique challenges posed by COVID-19.

OBJECTIVES

The ERO PRR is designed to establish requirements, as well as best practices, for all detention facilities housing ICE detainees to follow during the COVID-19 pandemic. Consistent with ICE detention standards, all facilities housing ICE detainees are required to have a COVID-19 mitigation plan that meets the following four objectives:

³ See, e.g., Attachment B, Centers of Disease Control and Prevention, *Symptoms of Coronavirus*, <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html> (last visited June 21, 2020).

⁴ See, e.g., Attachment C, ICE National Detention Standards 2019 (NDS), Standard 4.3, Medical Care, at II.D.2 (p. 114), https://www.ice.gov/doclib/detention-standards/2019/4_3.pdf; Attachment D, 2011 ICE Performance-Based National Detention Standards (PBNDS), Revised 2016, Standard 4.3, Part V.C.1 (p. 261), <https://www.ice.gov/doclib/detention-standards/2011/4-3.pdf>; Attachment E, 2008 ICE PBNDS, Standard 4-22, Medical Care, V.C.1 (pp. 5-6), https://www.ice.gov/doclib/dro/detention-standards/pdf/medical_care.pdf.

- To protect employees, contractors, detainees, visitors, and stakeholders from exposure to the virus;
- To maintain essential functions and services at the facility throughout the pendency of the pandemic;
- To reduce movement and limit interaction of detainees with others outside their assigned housing units, as well as staff and others, and to promote social distancing within housing units; and
- To establish means to monitor, cohort, quarantine, and isolate the sick from the well.⁵

COMPLIANCE MEASURES

To ensure that detention facilities comply with the detention requirements set forth in the ERO PRR, ICE will conduct bi-weekly spot checks at over 72-hour ICE detention facilities during the COVID-19 pandemic. Upon identification of a deficiency, ICE will provide written notice to the facility and allow seven business days for submission of a corrective action plan to ICE for approval. Life/safety issues identified by ICE will be corrected during the COVID-19 spot checks, if possible, or the facility will be required to submit a corrective action plan within three business days.

- For dedicated ICE detention facilities, which operate under Quality Assurance Surveillance Plans, ICE will issue a Contract Discrepancy Report (CDR), which may include contract sanctions, for failure to bring the facility into compliance with the minimum requirements of the ERO PRR within the ICE-approved timeframe. The CDR may become a part of the supporting documentation for contract payment deductions, fixed fee deductions, award fee nonpayment, or other contractual actions deemed necessary by the Contracting Officer. If the detention facility continues to have deficiencies despite the issuance of CDRs, ICE may seek to terminate the contract and/or decline to renew the contract.
- For non-dedicated ICE detention facilities that fail to meet the minimum requirements of the ERO PRR, ICE will issue a Notice of Intent indicating that the intergovernmental service agreement is in jeopardy due to non-compliance

⁵ A *cohort* is a group of persons with a similar condition grouped or housed together for observation over a period of time. Isolation and quarantine are public health practices used to protect the public from exposure to individuals who have or may have a contagious disease. For purposes of this document, and as defined by the CDC, *quarantine* is the separation of a person or group of people reasonably believed to have been exposed to a communicable disease but not yet symptomatic, from others who have not been exposed, to prevent the possible spread of the communicable disease. For purposes of this document, and as defined by the CDC, *isolation* is the separation of a person or group of people known or reasonably believed to be infected with a communicable disease and potentially infectious from others to prevent the spread of the communicable disease.

with the ERO PRR and may take appropriate action, including removing its detention population from the facility or reducing its detention population at the facility on a temporary or permanent basis, depending on the nature of the non-compliance.

CONCEPT OF OPERATIONS

The ERO PRR is intended for use across ICE's entire detention network, applying to all facilities housing ICE detainees, including ICE-owned Service Processing Centers, facilities operated by private vendors, and facilities operated by local government agencies that have mixed populations of which ICE detainees comprise only a small fraction.

DEDICATED ICE DETENTION FACILITIES

All dedicated ICE detention facilities⁶ must:

- Comply with the provisions of their relevant ICE contract or service agreement.
- Comply with the ICE national detention standards applicable to the facility, generally [PBNDS 2011](#).
- Comply with the CDC's [Interim Guidance on Management of Coronavirus Disease 2019 \(COVID-19\) in Correctional and Detention Facilities \(Attachment F\)](#).
- Follow ICE's March 27, 2020 Memorandum to Detention Wardens and Superintendents on COVID-19 Action Plan Revision 1, and subsequent updates (Attachment G).
- Report all confirmed and suspected COVID-19 cases to the local ERO Field Office Director (or designee), Field Medical Coordinator, and local health department immediately.
- Evaluate all new intakes within five days of entering ICE custody to determine whether the detainee falls within the populations identified by the CDC as potentially being at higher risk for serious illness from COVID-19, and/or the subclasses certified in *Fraihat v. ICE*, --- F. Supp. 3d ---, 2020 WL 1932570 (C.D. Cal. Apr. 20, 2020), and notify both the local ERO Field Office Director

⁶ Dedicated ICE detention facilities are facilities that house only ICE detainees. Dedicated ICE detention facilities may be ICE-owned Service Processing Centers, privately owned Contract Detention Facilities, or facilities operated by state or local governments that hold no other detention populations except ICE detainees.

(or designee) and the Field Medical Coordinator as soon as practicable, but in no case more than twelve hours of determining whether the detainee meets the criteria. These populations and subclasses include:

- People aged 55 and older;
- People who are pregnant;
- People of all ages with chronic health conditions, including:
 - Cardiovascular disease (congestive heart failure, history of myocardial infarction, history of cardiac surgery);
 - High blood pressure;
 - Chronic respiratory disease (asthma, chronic obstructive pulmonary disease including chronic bronchitis or emphysema, or other pulmonary diseases);
 - Diabetes;
 - Cancer;
 - Liver disease;
 - Kidney disease;
 - Autoimmune diseases (psoriasis, rheumatoid arthritis, systemic lupus erythematosus);
 - Severe psychiatric illness (including Psychotic Disorder, Bipolar Disorder, Schizophrenia or Schizoaffective Disorder, Major Depressive Disorder with Psychotic Features, Dementia and/or a Neurocognitive Disorder, or Intellectual Development Disorder (moderate, severe, or profound));
 - History of transplantation;
 - HIV/AIDS;
 - Severe obesity (body mass index [BMI] of 40 or higher).
- Individuals of all ages who are detained with a physical or mental impairment that substantially limits one or more major life activities or who has a record of physical or mental impairment that substantially limits a major life activity.

- Detainees who claim to meet the criteria should be evaluated within five days of making the claim and, upon evaluation, both the local ERO Field Office Director (or designee) and the Field Medical Coordinator must be notified as soon as practicable, but in no case more than twelve hours after the evaluation has occurred, as to whether the detainee meets the criteria.
- Notification shall be made via email from the facility's Health Services Administrator (HSA) (or equivalent) and contain the following subject line for ease of identification: "Notification of COVID-19 High Risk Detainee (A-Number)." At a minimum, the HSA email message will provide the following information:
 - Detainee name;
 - Detention location;
 - Current medical issues as well as medications currently prescribed;
 - Facility medical Point of Contact (POC) and phone number.

NON-DEDICATED ICE DETENTION FACILITIES

All non-dedicated ICE detention facilities and local jails housing ICE detainees must:

- Comply with the provisions of their relevant ICE contract or service agreement.
- Comply with the ICE national detention standards applicable to the facility, generally [PBNDS 2011](#).
- Comply with the [CDC *Interim Guidance on Management of Coronavirus Disease 2019 \(COVID-19\) in Correctional and Detention Facilities*](#).
- Report all confirmed and suspected COVID-19 cases to the local ERO Field Office Director (or designee), Field Medical Coordinator, and local health department immediately.
- Notify both the ERO Field Office Director (or designee) and Field Medical Coordinator as soon as practicable, but in no case more than twelve hours after identifying any detainee who meets the CDC's identified populations potentially being at higher risk for serious illness from COVID-19, and/or the subclasses certified in *Fraihat v. ICE, supra*. These populations and subclasses include:
 - People aged 55 and older;

- People who are pregnant;
 - People of all ages with chronic health conditions, including:
 - Cardiovascular disease (congestive heart failure, history of myocardial infarction, history of cardiac surgery);
 - High blood pressure;
 - Chronic respiratory disease (asthma, chronic obstructive pulmonary disease including chronic bronchitis or emphysema, or other pulmonary diseases);
 - Diabetes;
 - Cancer;
 - Liver disease;
 - Kidney disease;
 - Autoimmune diseases (psoriasis, rheumatoid arthritis, systemic lupus erythematosus);
 - Severe psychiatric illness (including Psychotic Disorder, Bipolar Disorder, Schizophrenia or Schizoaffective Disorder, Major Depressive Disorder with Psychotic Features, Dementia and/or a Neurocognitive Disorder, or Intellectual Development Disorder (moderate, severe, or profound));
 - History of transplantation;
 - HIV/AIDS;
 - Severe obesity (body mass index [BMI] of 40 or higher).
 - People of all ages who are detained with a physical or mental impairment that substantially limits one or more major life activities or who has a record of physical or mental impairment that substantially limits a major life activity.
- Notification should be made via email from the facility’s HSA (or equivalent) and should contain the following subject line for ease of identification: “Notification of COVID-19 High Risk Detainee (A-Number).” Other standardized means of communicating this information to ICE, as established by

agreement between the local ERO Field Office Director (or designee) and the Warden or Superintendent, are acceptable. At a minimum the HSA communication to ICE will provide the following information:

- Detainee name;
- Detention location;
- Current medical issues as well as medications currently prescribed;
- Facility medical POC and phone number.

ALL FACILITIES HOUSING ICE DETAINEES

In addition to the specific measures listed above, all detention facilities housing ICE detainees must also comply with the following:

PREPAREDNESS

Administrators should plan and prepare for COVID-19 by ensuring that all persons in the facility know the symptoms of COVID-19 and how to respond if they develop symptoms. Other essential actions include developing contingency plans for reduced workforces due to absences, coordinating with public health and correctional partners, and communicating clearly with staff and detainees about these preparations and how they may temporarily alter daily life.

- Develop information-sharing systems with partners.
 - Identify points of contact in relevant state, local, tribal, and/or territorial public health department before cases develop.
 - Communicate with other correctional and detention facilities in the same geographic area to share information including disease surveillance and absenteeism patterns among staff.
- Review existing pandemic, influenza, all-hazards, and disaster plans, and revise for COVID-19, and ensure that they meet the requirements of ICE's detention standards.
- Offer the seasonal influenza vaccine to all detained persons (existing populations and new intakes) and staff throughout the influenza season, where possible.
- Staffing
 - Review sick leave policies to ensure that staff can stay home when sick and determine which officials will have the authority to send symptomatic staff

home. Staff who report for work with symptoms of COVID-19 must be sent home and advised to follow CDC-recommended steps for persons exhibiting COVID-19 symptoms.

- Staff who test positive for COVID-19 must inform their workplace and personal contacts immediately. If a staff member has a confirmed COVID-19 infection, the relevant employers will inform other staff of their possible exposure to COVID-19 in the workplace consistent with any legal limitations on the sharing of such information. Exposed employees must then self-monitor for symptoms (e.g., fever, cough, or shortness of breath).
- Identify staff whose duties would allow them to work from home in order to promote social distancing and further reduce the risk of COVID-19 transmission.
- Determine minimum levels of staff in all categories required for the facility to function safely.
- Follow the Public Health Recommendations for Community-Related Exposure.⁷

➤ Supplies

- Ensure that sufficient stocks of hygiene supplies (soap, hand sanitizer, tissues), personal protective equipment (PPE) (facemasks, N95 respirators, eye protection, disposable medical gloves, and disposable gowns/one-piece coveralls), and medical supplies (consistent with the healthcare capabilities of the facility) are on hand and have a plan in place to restock as needed if COVID-19 transmission occurs within the facility.
- Note that shortages of N95 respirators are anticipated during the COVID-19 response. Based on local and regional situational analysis of PPE supplies, face masks should be used when the supply chain of N95 respirators cannot meet the demand.
- Follow COVID-19: Strategies for Optimizing the Supply of PPE.⁸
- Soiled PPE items should be disposed in leak-proof plastic bags that are tied at the top and not re-opened. Bags can be disposed of in the regular solid waste

⁷ Attachment H, Centers of Disease Control and Prevention, *Public Health Guidance for Community-Related Exposure*, <https://www.cdc.gov/coronavirus/2019-ncov/php/public-health-recommendations.html> (last visited June 21, 2020).

⁸ Attachment I, Centers for Disease Control and Prevention, *Strategies to Optimize the Supply of PPE and Equipment*, <https://www.cdc.gov/coronavirus/2019-ncov/hcp/ppe-strategy/> (last visited June 21 2020).

stream.

- Cloth face coverings should be worn by detainees and staff (when PPE supply is limited) to help slow the spread of COVID-19. Cloth face masks should:
 - Fit snugly but comfortably against the side of the face be secured with ties or ear loops where possible or securely tied;
 - Include multiple layers of fabric;
 - Allow for breathing without restriction;
 - Be able to be laundered and machine dried without damage or change to shape.

➤ Hygiene

- Reinforce healthy hygiene practices and provide and restock hygiene supplies throughout the facility, including in bathrooms, food preparation and dining areas, intake areas, visitor entries and exits, visitation rooms, common areas, medical, and staff-restricted areas (e.g., break rooms).
- Require all persons within the facility to cover their mouths and noses with their elbows (or ideally with a tissue) rather than with their hands when they cough or sneeze, and to throw all tissues in the trash immediately after use.
- Provide detainees and staff no-cost access to tissues and no-touch receptacles for disposal.
- Require all persons within the facility to maintain good hand hygiene by regularly washing their hands with soap and water for at least 20 seconds, especially after coughing, sneezing, or blowing their noses; after using the bathroom; before eating or preparing food; before taking medication; and after touching garbage.
- Provide detainees and staff no-cost, unlimited access to supplies for hand cleansing, including liquid soap, running water, hand drying machines or disposable paper towels, and no-touch trash receptacles.
- Provide alcohol-based hand sanitizer with at least 60 percent alcohol where permissible based on security restrictions.
- Require all persons within the facility to avoid touching their eyes, noses, or mouths without cleaning their hands first.
- Post signage throughout the facility reminding detained persons and staff to

practice good hand hygiene and cough etiquette (printable materials for community-based settings can be found on the [CDC website](#)). Signage must be in English and Spanish, as well as any other common languages for the detainee population at the facility.

- Prohibit sharing of eating utensils, dishes, and cups.
- Prohibit non-essential personal contact such as handshakes, hugs, and high-fives.

➤ **Cleaning/Disinfecting Practices**

- Adhere to CDC recommendations for cleaning and disinfection during the COVID-19 response.⁹
- Several times a day using household cleaners and Environmental Protection Agency-registered disinfectants, clean and disinfect surfaces and objects that are frequently touched, especially in common areas (e.g., doorknobs, light switches, sink handles, countertops, toilets, toilet handles, recreation equipment). The Environmental Protection Agency's (EPA) list of certified cleaning products is located here.
- Staff should clean shared equipment several times per day and on a conclusion-of-use basis (e.g., radios, service weapons, keys, handcuffs).
- Ensure that transport vehicles are thoroughly cleaned after carrying a confirmed or suspected COVID-19 case.
- Facility leadership will ensure that there is adequate oversight and supervision of all individuals responsible for cleaning and disinfecting these areas.

CDC Recommended Cleaning Tips

➤ **Hard (Non-porous) Surfaces**

- If surfaces are dirty, they should be cleaned using a detergent or soap and water prior to disinfection.
- For disinfection, most common EPA-registered household disinfectants should be effective.
 - A list of products that are EPA-approved for use against the virus that

⁹ Attachment J, Centers for Disease Control and Prevention, Cleaning and Disinfection for Community Facilities, <https://www.cdc.gov/coronavirus/2019-ncov/community/organizations/cleaning-disinfection.html> (last visited June 212020).

causes COVID-19 is available [here](#). Follow the manufacturer's instructions for all cleaning and disinfection products for concentration, application method and contact time, etc.

- Additionally, diluted household bleach solutions (at least 1000 ppm sodium hypochlorite) can be used if appropriate for the surface. Follow manufacturer's instructions for application, ensuring a contact time of at least one minute and allowing proper ventilation during and after application. Check to ensure the product is not past its expiration date. Never mix household bleach with ammonia or any other cleanser. Unexpired household bleach will be effective against coronaviruses when properly diluted.
 - Prepare a bleach solution by mixing:
 - 5 tablespoons (1/3 cup) bleach per gallon of water; or
 - 4 teaspoons bleach per quart of water.

➤ Soft (Porous) Surfaces

- For soft (porous) surfaces such as carpeted floor, rugs, and drapes, remove visible contamination if present and clean with appropriate cleaners indicated for use on these surfaces. After cleaning:
 - If the items can be laundered, launder items in accordance with the manufacturer's instructions using the warmest appropriate water setting for the items and then dry items completely.
 - Otherwise, use products that are EPA-approved for use against the virus that causes COVID-19 and that are suitable for porous surfaces.¹⁰

➤ Electronics

- For electronics such as tablets, touch screens, keyboards, remote controls, and ATM machines, remove visible contamination if present.
 - Follow the manufacturer's instructions for all cleaning and disinfection products.
 - Consider use of wipeable covers for electronics.

¹⁰ Attachment K, U.S. Environmental Protection Agency, *List N: Disinfectants for Use Against SARS-CoV-2*, <https://www.epa.gov/pesticide-registration/list-n-disinfectants-use-against-sars-cov-2> (last visited June 21, 2020).

- If no manufacturer guidance is available, consider the use of alcohol-based wipes or sprays containing at least 70% alcohol to disinfect touch screens. Dry surfaces thoroughly to avoid pooling of liquids.
- Linens, Clothing, and Other Items That Go in the Laundry
 - In order to minimize the possibility of dispersing virus through the air, do not shake dirty laundry.
 - Wash items as appropriate in accordance with the manufacturer’s instructions. If possible, launder items using the warmest appropriate water setting for the items and dry items completely. Dirty laundry that has been in contact with an ill person can be washed with other people’s items.
 - Clean and disinfect hampers or other carts for transporting laundry according to guidance above for hard or soft surfaces.

PREVENTION

Detention facilities can prevent the introduction of COVID-19 from the community and reduce transmission if it is already inside by reinforcing good hygiene practices among incarcerated/detained persons, staff, and visitors (including increasing access to soap and paper towels), intensifying cleaning/disinfection practices, and implementing social distancing strategies.

Because many individuals infected with COVID-19 do not display symptoms, the virus could be present in facilities before cases are identified. Both good hygiene practices and social distancing are critical in preventing further transmission.

- Perform pre-intake screening for all staff and new entrants for symptoms of COVID-19.

Screening should take place before staff and new intakes enter the facility or just inside the facility, where practicable. For new admissions, this should occur before beginning the intake process, in order to identify and immediately isolate any detainee with symptoms before the individual comes in contact with others or is placed in the general population. This should include temperature screening of all staff and new entrants, as well as a verbal symptoms check.

- Verbal screening for symptoms of COVID-19 and contact with COVID-19 cases should include the following questions based on [Interim Guidance: Managing COVID-19 in Correctional/Detention Facilities](#):
 - Today or in the past 24 hours, have you had any of the following symptoms:

- Fever, felt feverish, or had chills?
 - Cough?
 - Difficulty breathing?
 - Chills?
 - Muscle pain?
 - Sore throat?
 - New loss of taste or smell?
- In the past fourteen days, have you had contact with a person known to be infected with COVID-19 where you were not wearing the recommended proper PPE?
- If staff have symptoms of COVID-19 (e.g., fever, cough, shortness of breath), they must be denied access to the facility.
- If any new intake has symptoms of COVID-19:
 - Require the individual to wear a face mask.
 - Ensure that staff interacting with the symptomatic individual wears recommended PPE.
 - Isolate the individual and refer to healthcare staff for further evaluation.
 - Facilities without onsite healthcare staff should contact their state, local, tribal, and/or territorial health department to coordinate effective isolation and necessary medical care.
- If an individual is a close contact of a known COVID-19 case or has traveled to an affected area, but has no COVID-19 symptoms, quarantine the individual and monitor for symptoms two times per day for fourteen days.

➤ Visitation

- During suspended (social) or modified (legal) visitation programs, the facilities should provide access to virtual visitation options where available. When not possible, verbally screen all visitors on entry for symptoms of COVID-19 and perform temperature checks, when possible. ICE continues to explore opportunities to enhance attorney access while legal visits are being impacted. For facilities at which immigration hearings are conducted or where detainees are otherwise held who have cases pending immigration

proceedings, this may include

- Adding all immigration attorneys of record to the Talton Pro-bono platform.
 - Requiring facilities to establish a process for detainees/attorneys to schedule appointments and facilitate the calls.
 - Leveraging technology (e.g., tablets, smartphones) to facilitate attorney/client communication.
 - Working with the various detention contractors and telephone service providers to ensure that all detainees receive some number of free calls per week.
 - Communicate with the public about any changes to facility operations, including visitation programs. Facilities are encouraged to prohibit or, at a minimum, significantly adopt restricted visitation programs, and are required to suspend all volunteer work program (VWP) assignments for detainees assigned to food service and other VWP assignments, where applicable, that require individuals to interact with each other at distances of less than six feet. Any detainee participating in a VWP assignment is required to wear appropriate PPE for the position at all times (e.g., disposable gloves, masks, goggles). Detainees in isolation or quarantine may not be assigned to a VWP detail.
- Where possible, restrict transfers of detained non-ICE populations to and from other jurisdictions and facilities unless necessary for medical evaluation, isolation/quarantine, clinical care, or extenuating security concerns.
 - Consider suspending work release programs for inmates at shared facilities to reduce overall risk of introduction and transmission of COVID-19 into the facility.
 - Require all staff (both medical and correctional) to wear PPE when encountering or interacting with any ICE detainee at a distance of less than six feet.
 - Required PPE should always be worn by staff, even if separated by a distance of six feet or more, if the individual appears feverish or ill and/or with respiratory symptoms while interviewing, escorting, or interacting in other ways.
 - Additional Measures to Facilitate Social Distancing
 - Although strict social distancing may not be possible in congregate settings such as detention facilities, all facilities housing ICE detainees should implement the following measures to the extent practicable:

- Efforts should be made to reduce the population to approximately 75 percent of capacity.
- Where detainee populations are such that cells are available, to the extent possible, house detainees in individual rooms.
- Recommend that detainees sharing sleeping quarters sleep “head-to-foot.”
- Extend recreation, law library, and meal hours and stagger detainee access to the same in order to limit the number of interactions between detainees from other housing units.
- Staff and detainees should be directed to avoid congregating in groups of ten or more, employing social distancing strategies at all times.
- Whenever possible, all staff and detainees should maintain a distance of six feet from one another.
- If practicable, beds in housing units should be rearranged to allow for six feet of distance between the faces of detainees.

MANAGEMENT

If there has been a suspected COVID-19 case inside the facility (among incarcerated/detained persons, staff, or visitors who have recently been inside), begin implementing management strategies while test results are pending. Essential management strategies include placing cases and individuals with symptoms under medical isolation, quarantining their close contacts, and facilitating necessary medical care, while observing relevant infection control and environmental disinfection protocols and wearing recommended PPE.

ICE Custody Review for Potentially High-Risk Detainees

Upon being informed of a detainee who may potentially be at higher risk for serious illness from exposure to COVID-19, ERO will review the case to determine whether continued detention is appropriate.¹¹ ICE will make such custody determinations on a case-by-case basis, pursuant to the applicable legal standards, with due consideration of the public health considerations implicated.

- Considerable effort should be made to quarantine all new entrants for fourteen days before they enter the general population.
- To do this, facilities should consider cohorting daily intakes; two days of new intakes, or multiple days of new intakes, in designated areas prior to placement into

¹¹ Attachment L, Assistant Director Peter Berg, Enforcement and Removal Operations, *Updated Guidance: COVID-19 Detained Docket Review* (Apr. 4, 2020).

the general population. Given significant variations among facilities, cohorting options and capabilities will differ across ICE's detention network. ICE encourages all facilities to adopt the most effective cohorting methods practicable based on the individual facility characteristics taking into account the number of new intakes anticipated per day.

- For suspected or confirmed COVID-19 cases:
 - Isolate the individual immediately in a separate environment from other individuals. Facilities should make every possible effort to isolate persons individually. Each isolated individual should be assigned his or her own housing space and bathroom where possible. Isolating ill detainees as a group should only be practiced if there are no other available options.
 - If single isolation rooms are unavailable, individuals with laboratory-confirmed COVID-19 should be isolated together as a cohort separate from other detainees, including those with pending tests. Ill detainees that are pending testing or are waiting for test results should be isolated together as a group separate from laboratory-confirmed COVID-19 cases and other detainees. Confirmed COVID-19 cases should not be cohorted with suspected cases or case contacts.
 - Housing should maintain separation of groups by common criteria (e.g., COVID-19 test results positive, febrile or symptomatic pending testing or results, asymptomatic/exposed).
 - Ensure that the individual is always wearing a face mask (if it does not restrict breathing) when outside of the isolation space, and whenever another individual enters the isolation room. If wearing masks will negatively impact breathing, facilities should ensure caregivers are aware of that fact and implement restrictions on contact as appropriate during isolation (e.g., increased social distancing, PPE use by people who enter space, moving and handling people separately, increased cleaning, etc.). Masks should be changed at least daily, and when visibly soiled or wet.
 - In the event that a facility requires more isolation beds for detainees, ICE must be promptly notified so that transfers to other facilities, transfers to hospitals, or releases can be coordinated immediately. Until such time as the transfer or release is arranged, the facility must be especially mindful of cases that are at higher risk of severe illness from COVID-19. Ideally, ill detainees should not be isolated with other infected individuals. If isolating of ill detainees as a group is unavoidable, make all possible accommodations until transfer occurs to prevent transmission of other infectious diseases to the higher-risk individual (e.g., allocate more space for a higher-risk individual within a shared isolation room).

- Review the CDC’s preferred method of medically isolating COVID-19 cases [here](#), depending on the space available in a particular facility. In order of preference, individuals under medical isolation should be housed:
 - Separately, in single cells with solid walls (i.e., not bars) and solid doors that close fully.
 - Separately, in single cells with solid walls but without solid doors.
 - As a cohort, in a large, well-ventilated cell with solid walls and a solid door that closes fully. Employ social distancing strategies related to housing in the Prevention section above.
 - As a cohort, in a large, well-ventilated cell with solid walls but without a solid door. Employ social distancing strategies related to housing in the Prevention section above.
 - As a cohort, in single cells without solid walls or solid doors (i.e., cells enclosed entirely with bars), preferably with an empty cell between occupied cells. (Although individuals are in single cells in this scenario, the airflow between cells essentially makes it a cohort arrangement in the context of COVID-19.)
 - As a cohort, in multi-person cells without solid walls or solid doors (i.e., cells enclosed entirely with bars), preferably with an empty cell between occupied cells. Employ social distancing strategies related to housing in the Prevention section above.
- Maintain isolation until all the CDC criteria have been met. The decision to discontinue Transmission-Based Precautions should be made using a test-based strategy, a symptom-based strategy, or a time-based strategy.¹² Detainees that meet criteria for discontinuation of transmission-based precautions can be housed in general population.
 - Symptom-based strategy: Patients initially presenting with fever and/or other symptoms
 - At least three days (72 hours) have passed since recovery defined as resolution of fever without the use of fever-reducing medications and improvement in respiratory symptoms (e.g., cough, shortness of breath);
 - At least seven days have passed since symptoms first appeared;

¹² See Attachment M, Centers for Disease Control and Prevention, *Discontinuation of Transmission-Based Precautions and Disposition of Patients with COVID-19 in Healthcare Settings (Interim Guidance)*, <https://www.cdc.gov/coronavirus/2019-ncov/hcp/disposition-hospitalized-patients.html> (last visited June 21, 2020).

and

- A minimum of fourteen days has passed since the first specimen collection resulting in a positive confirmatory test for a U.S. Food and Drug Administration (FDA) Emergency Use Authorization (EUA) COVID-19 molecular assay for detection of SARS-CoV-2.
- Time-based strategy: Patients initially testing positive and presenting without fever or symptoms
 - A minimum of fourteen days has passed since the first specimen collection resulting in a positive confirmatory test for an FDA EUA COVID-19 molecular assay for detection of SARS-CoV-2;
 - If the detainee develops signs and/or symptoms consistent with COVID-19, the release from isolation precautions needs to be determined using the symptom-based strategy above.
- Test-based strategy: Patients initially presenting with or without fever and/or symptoms (preferred if the patient is immunocompromised)
 - Resolution of fever without the use of fever-reducing medications (if initially febrile);
 - in symptoms (if initially present);
 - Negative results of an FDA EUA COVID-19 molecular assay for detection of SARS-CoV-2 from at least two consecutive respiratory specimens collected at least 24 hours apart (total of two negative specimens),¹³ and
 - A minimum of fourteen days has passed since the first specimen collection resulting in a positive confirmatory test with an FDA EUA COVID-19 molecular assay for detection of SARS-CoV-2.
- Meals should be provided to COVID-19 cases in their isolation rooms. Isolated cases should throw disposable food service items in the trash in their isolation room. Non-disposable food service items should be handled with gloves and washed with hot water or in a dishwasher. Individuals handling used food service items must clean their hands after removing gloves.
- Laundry from a COVID-19 case can be washed with other individuals' laundry.

¹³Attachment N, Centers of Disease Control and Prevention, *Interim Guidelines for Collecting, Handling, and Testing Clinical Specimens for COVID-19*, <https://www.cdc.gov/coronavirus/2019-ncov/lab/guidelines-clinical-specimens.html> (last visited June 21, 2020).

- Individuals handling laundry from COVID-19 cases should wear disposable gloves, discard gloves after each use, and clean their hands after handling.
 - Do not shake dirty laundry. This will minimize the possibility of dispersing the virus through the air.
 - Launder items as appropriate in accordance with the manufacturer’s instructions. If possible, launder items using the warmest appropriate water setting for the items and dry items completely.
 - Clean and disinfect clothes hampers according to guidance above for surfaces. If permissible, consider using a bag liner that either is disposable or can be laundered.
- Consistent with CDC recommendations, facilities “considering diagnostic testing of people with possible COVID-19 should continue to work with their local and state health departments to coordinate testing through public health laboratories, or work with commercial or clinical laboratories using diagnostic tests authorized for emergency use by the U.S. Food and Drug Administration.”¹⁴
- For CDC recommendations for antibody testing, please go to: <https://www.cdc.gov/coronavirus/2019-ncov/lab/resources/antibody-tests.html>
 - For CDC recommendations for viral testing and specimen collection, please go to: <https://www.cdc.gov/coronavirus/2019-nCoV/hcp/clinical-criteria.html>

¹⁴Attachment O, Centers of Disease Control and Prevention, *Overview of Testing for SARS-CoV-2*, <https://www.cdc.gov/coronavirus/2019-nCoV/hcp/clinical-criteria.html> (last visited June 21, 2020).

ATTACHMENTS

ATTACHMENT LETTER	DOCUMENT NAME AND CITATION
A	U.S. Immigration and Customs Enforcement, <i>Updated ICE statement on COVID-19</i> (Mar. 18, 2020), https://www.ice.gov/news/releases/updated-ice-statement-covid-19 .
B	Centers of Disease Control and Prevention, <i>Symptoms of Coronavirus</i> , https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html (last visited June 21, 2020).
C	ICE National Detention Standards 2019, Standard 4.3, Medical Care, https://www.ice.gov/doclib/detention-standards/2019/4_3.pdf .
D	2011 ICE Performance-Based National Detention Standards, Revised 2016, Standard 4.3, https://www.ice.gov/doclib/detention-standards/2011/4-3.pdf .
E	2008 ICE Performance-Based National Detention Standards, Standard 4-22, Medical Care, https://www.ice.gov/doclib/dro/detention-standards/pdf/medical_care.pdf .
F	Centers of Disease Control and Prevention, <i>Interim Guidance on Management of Coronavirus Disease 2019 (COVID-19) in Correctional and Detention Facilities</i> (Mar. 23, 2020), https://www.cdc.gov/coronavirus/2019-ncov/downloads/guidance-correctional-detention.pdf (last visited June 21, 2020).
G	Memorandum from Executive Associate Director Enrique Lucero, Enforcement and Removal Operations, <i>Memorandum on Coronavirus 2019 (COVID-19) Action Plan, Revision 1</i> (Mar. 27, 2020).
H	Centers of Disease Control and Prevention, <i>Public Health Guidance for Community-Related Exposure</i> , https://www.cdc.gov/coronavirus/2019-ncov/php/public-health-recommendations.html (last visited June 21, 2020).
I	Centers for Disease Control and Prevention, <i>Strategies to Optimize the Supply of PPE and Equipment</i> , https://www.cdc.gov/coronavirus/2019-ncov/hcp/ppe-strategy/ (last visited June 21, 2020).

ATTACHMENT LETTER	DOCUMENT NAME AND CITATION
J	Centers for Disease Control and Prevention, <i>Cleaning and Disinfection for Community Facilities</i> , https://www.cdc.gov/coronavirus/2019-ncov/community/organizations/cleaning-disinfection.html (last visited June 21, 2020).
K	U.S. Environmental Protection Agency, <i>List N: Disinfectants for Use Against SARS-CoV-2</i> , https://www.epa.gov/pesticide-registration/list-n-disinfectants-use-against-sars-cov-2 (last visited June 21, 2020).
L	Assistant Director Peter Berg, Enforcement and Removal Operations, <i>Updated Guidance: COVID-19 Detained Docket Review</i> (Apr. 4, 2020).
M	Centers for Disease Control and Prevention, <i>Discontinuation of Transmission-Based Precautions and Disposition of Patients with COVID-19 in Healthcare Settings (Interim Guidance)</i> , https://www.cdc.gov/coronavirus/2019-ncov/hcp/disposition-hospitalized-patients.html (last visited June 21, 2020)
N	Centers of Disease Control and Prevention, <i>Interim Guidelines for Collecting, Handling, and Testing Clinical Specimens for COVID-19</i> , https://www.cdc.gov/coronavirus/2019-ncov/lab/guidelines-clinical-specimens.html (last visited June 21, 2020)
O	Centers of Disease Control and Prevention, <i>Overview of Testing for SARS-CoV-2</i> , https://www.cdc.gov/coronavirus/2019-nCoV/hcp/clinical-criteria.html (last visited June 21, 2020)